

Morpho-physiological Effects of Drought on Medicinal Plants and the Potential Use of Remote Sensing - A Review

Tewfik Al-hamed^{1*}, Safwan M. Shiyab², and Jawad Al-Bakri³

¹Department of Horticulture and Crop Science, Faculty of Agriculture, University of Jordan.

²Department of Horticulture and Crop Science, Faculty of Agriculture, University of Jordan.

³Department of Land, Water, and Environment, Faculty of Agriculture, University of Jordan.

Received 9 August 2021; Accepted 13 January 2022

Abstract

This review addresses the growth and development of medicinal plants under arid land conditions and the potential use of remote sensing technologies to map their distribution, as well as morphological and physiological responses in Arid lands. Plant morpho-physiological variables including, chlorophyll content, and gas exchange (photosynthesis, transpiration, vapor pressure, and stomatal conductance) are linked to plant water status. Multispectral and hyperspectral remote sensing techniques are promising for detecting morphological physiological changes. Vegetation indices derived from hyperspectral and multispectral imagery makes it possible to assess medicinal plants' health through successful detection of the ground plant physiological variables and canopy cover including chlorophyll, canopy density, gas exchange (red, 600-700 nm, near-infrared spectrum, 700-1100 nm) as well as water status (shortwave infrared, 1300-2500 nm). Surface reflectance data within the shortwave infrared bands (water bands) revealed significant differences between well-watered and drought-stressed plants. However, the moderate spatial resolution (Sentinel-2: 20 m, Landsat: 30 m) for the space-born free sensors and the need for a cloud-free sky could be limitations. Overall, vegetation indices derived from remotely sensed data are a useful approach for estimating the physiology of plants (medicinal plants) especially those under drought stress.

© 2022 Jordan Journal of Earth and Environmental Sciences. All rights reserved

Keywords: Medicinal Plants, Drought Stress, Water Status, Hyperspectral, Multispectral, Vegetation Index.

1. Introduction

Climate change is one of the major problems facing the world because it is foretelling climate patterns changes and a higher frequency of risky weather events. In recent years, the frequency and severity of droughts in the Mediterranean region have increased due to global warming (IPCC, 2013; Trenberth et al., 2014). Generally, climate change has a potential impact on agriculture, coastal areas, biodiversity, urban systems, society, water, and health sectors (FAO, 2008). In the agricultural sector, climate-change-related risks are represented by higher temperatures, rainfall decreases, the shift in the rainy season and seasonal alterations, heatwaves, and extreme events especially heavy rainfall and droughts (WB, 2021). Drought is a disastrous natural phenomenon that has negative impacts on plants and especially under the changing climate, it becomes more frequent and severe (Seneviratne et al., 2012). Drought has destructive effects on socioeconomic, plants, and the environment because it leads to insufficient precipitation, high evapotranspiration, and over-exploitation of water resources (Yurekli and Kurunc, 2004; Bhuiyan et al., 2006; García-Caparrós et al., 2019). Drought effects can be classified as direct and indirect (Van Lanen and Peters, 2000). Direct impacts of drought are the results of interactions among water deficiencies and environmental, and socio-economical components while indirect are a secondary result of water deficiency and are often occurred far away from the drought-impacted region (UNDRR, 2021).

Generally, drought can be categorized into three basic types in terms of measuring drought as a physical phenomenon; meteorological, agricultural, or hydrological drought (Wilhite and Glantz, 1985). Meteorological drought is a prolonged abnormal dryness period (compared to normal average precipitation), agricultural drought links meteorological variables (e.g., precipitation) to agricultural impacts such as soil water deficits while hydrological drought is associated with the influence of rainfall periods on the surface or groundwater supply (NDMC, 2021). In arid and semi-arid land, functional landscapes and their associated vegetation are ultimately dependent on water availability, which significantly affect vegetation distribution (Tardieu et al., 2018). Drought stress in plants (e.g., *Thymus citriodorus*) normally plants moisture content (dehydration) followed by a reduction in the metabolism process and photosynthesis and consequently plants death (Tátrai et al., 2016). However, the ability of plants to survive under stressed conditions depends on plant species, growth stage, duration, and the intensity of water deficit (Blum, 2017). Plant tolerance to abiotic stress such as water stress is unpredictable because of the complicated interactions between stress factors and plant molecular, biochemical, and physiological components associated with growth and development processes (Razmjoo et al., 2008).

Medicinal plants are the major sources of numerous valuable chemicals and/or drugs worldwide. According to the International Union for Conservation of Nature (IUCN)

* Corresponding author e-mail: Tewfik_Alhamad@mwi.gov.jo

and the World Wildlife Fund (WWF), about 50,000–80,000 flowering plants are used because of their medicinal values. In Europe, over 1300 medicinal plants are used, especially those from the wild sources which account for 90% of the total medicinal plants (Hao, 2019). Jordan is home to an abundance of flora and fauna, about 2978 plant species belonging to 120 families and 719 genera are recorded in Jordan (Abdelhalim et al., 2017). In addition, 20% of the total flora species in Jordan are classified as medicinal plants (Abdelhalim et al., 2017). High demand for medicinal plants coupled with unpredictable environmental stressors such as low rainfall has led to a significant reduction in their abundance worldwide. Cultural practices such as planting method and date, fertilizer application, irrigation regimes, and harvesting should be optimized to increase the growth and productivity of medicinal plants (Tanga et al., 2018). Efforts to conserve wildlife flora including medicinal plants have been observed recently including research studies focused on the growth (e.g soilless culture), physiology, production (yield, phenolic compounds, and oil content), and conservation of those plants (Al-Karaki and Othman 2009; Leskovar and Othman, 2016; Sharma et al., 2020).

Management and monitoring of wild plants including medicinal ones require frequent and consistent assessment of the canopy status and health over time. The main health indicators are morphology and physiological variables including chlorophyll contents, gas exchange [photosynthesis (P_n), transpiration (E), vapor pressure deficit (VPD), and stomatal conductance (g_s)] as well as plant development and yield (Tahat et al., 2020). These morpho-physiological variables can provide accurate information about the plant carbon assimilation rate, leaf water level (Othman et al., 2014a; Tadros et al., 2021), plant nutrition, fruit quality, and consequently their health (Al-Ajlouni et al., 2017; A'saf et al., 2020; Ayad et al., 2018; Alsmirat et al. 2018; Leskovar and Othman, 2018, 2021; Tahat et al., 2020). However, these methods of assessment are time-consuming, labor intensive, and expensive. In addition, the distribution of some wildlife flora could be not accessible. Another possible alternative to assess plant health is through remote sensing techniques (Al-Kofahi et al., 2019; Othman et al., 2014b, 2021; Tadros et al., 2020). Remote sensing has potentially assessed chlorophyll content (Othman et al., 2019), canopy density (Othman and St. Hilaire, 2021; Tadros et al., 2020), gas exchange and water status (Othman et al., 2014b, 2015), and geographic distribution (Al-Bakri et al., 2011). Remotely sensed data are providing an upscale view of the land and a spatiotemporal context to measure drought impacts (Hazaymeh and Hassan, 2017). In addition, vegetation indices derived from space-borne sensors (e.g., Landsat and MODIS) such as Normalized Difference Vegetation Index (NDVI) and the Vegetation Condition Index (VCI) have been successfully linked to chlorophyll content and water content in the plants (Rousta et al., 2020). In this review, we evaluated the usefulness of using the remote sensing approach as an alternative for evaluating medicinal plants' morpho-physiological responses under drought conditions.

2. Evaluation of Eco-physiological Parameters of Medicinal Plants

In the developing world, local communities depend on traditional medicine for primary health care (Jeelani et al. 2018). In addition, developed countries heavily use medicinal plants for their pharmaceutical products (Chapman and Chomchalow, 2005). Medicinal plants have several phenolic and antioxidant compounds which used to support human health and cure some diseases. In Jordan, several medicinal plant species are native to Eastern and Northern rangelands (Badia) (Oran and Al-Eisawi, 2015). More than 49 plant families and 120 plant species found in Jordan are used by neighborhood people for medical purposes (Atta and Alkofahi, 1998). However, uncontrolled grazing and frequent drought periods in the last decade reduced their numbers (Oran and Al-Eisawi, 2015). Therefore, research studies that focus on improving the tolerance of the medicinal plant to harsh conditions as well as controlling grazing intensity are essential for the potential sustainability of those plants.

Abiotic, and biotic (soil microorganisms) factors can significantly affect medicinal plant metabolite biosynthesis. Abiotic stresses such as temperature extremes and water stress can negatively affect the physiology and chemical composition of plants, which as a result induce abnormalities in medicinal plant metabolic processes such as growth, photosynthesis performance, and yield (Zaid et al., 2021). Optimal environmental conditions tend to increase medicinal plant biomass rather than synthesizing secondary metabolites (Pavarini et al., 2012). Postharvest processing can also lead to an irreversible quality loss of medicinal plants (Tanga et al., 2018). Therefore, finding proper management practices to guarantee high plant growth, productivity and quality are critical for the medicinal plants industry. In this context, management practices including irrigation, fertilization, and pest control required frequent assessment and evaluation during the growing season to sustain plant health and productivity. To assess the growth performance in response to those cultural practices (irrigation, fertilization, pest management) several plant-base and soil-base measurements have been recommended including water status (e.g. midday stem water potential (Ψ_{smd}), relative water content) and chlorophyll content (Othman et al., 2014b, 2015).

Plant-based measurements such as midday stem water potential, relative water content, gas exchange (P_n , g_s , E , VPD , respectively), and leaf pigments are viable approaches to assess the physiology of plants which depict their health status (Jones, 2004; Khasawneh et al., 2021; Leskovar and Othman, 2019; Othman and Leskovar, 2018, 2019). This is because many features of the tree's physiology react immediately to changes in tree tissues including water status (Jones, 2004). For example, Othman et al., (2014a) reported that under deficit water stress, Ψ_{smd} and gas exchange were significantly lower and resulted in lower growth rate and yield compared to non-water-stressed plants. Although total soluble sugars and proline concentration in chamomile (*Matricaria chamomilla* L.) was similar across water deficit regimes, the total chlorophyll concentration was significantly reduced in water deficit plants (55% of field capacity) compared to control (100% field capacity) (Pirzad et al., 2011).

Chlorophyll and carotenoids (carotenes and xanthophyll) are the major pigments of green leaves (Gitelson et al., 2002; Li et al., 2018). Pigment concentration inside the leaves potentially reflects the physiological performance of plants (Gamon and Surfus, 1999; Blackburn, 2007). This is because leaf pigments (chlorophyll, and anthocyanin) level in the leaves could act as a photoprotective mechanism that mitigates water deficit impact (Gamon and Surfus, 1999; Gori et al., 2021). Chlorophyll fluorescence (Fv/Fm) is a leaf-level physiological measurement that is used to study the performance of chlorophyll pigments under stress such as drought (Guidi et al., 2019). The decrease in Fv/Fm value is an indicator of low performance of photosynthetic pigments specifically, photosystem II (PSII) efficiency (Guidi et al., 2019). However, chlorophyll fluorescence measurements (re-emitted light from PSII) can interfere with the sunlight and thus many early systems had to be used in darkness and/or highly controlled light environments and it remains stable under mild and moderate drought stress and shows a significant decline only under severe drought conditions (Othman et al., 2014a). Gas exchange also is used in assessing plant performance including those studies focusing on plant growth and productivity as well as fruit quality (Leskovar and Othman, 2021; Kasaweneh et al., 2021). Gas exchange variables (Pn, gs, and E) normally decreased in parallel with increased stressors such as water deficit and nutrient deficiency (Ayad et al., 2018).

Water status is one of the most important factors limiting medicinal plant performance, especially those grown in an arid environment (Khorasaninejad et al., 2010; Pirzad et al., 2011). For example, drought stress significantly reduced the growth rate, essential oil, and yield of mint (*Mentha piperita* L.) compared to those grown under 100% field capacity (Khorasaninejad et al., 2010). Low soil moisture reduces stomatal conductance, total lipid content, photosynthetic capacity, chlorophyll content, transpiration, and the total dry matter (Guerfel et al., 2008; Grams et al., 2007; He et al., 2007; Warren et al., 2007; Damour et al., 2009; Arunyanark et al., 2008). At moderate drought, Pn decreased slightly because of the reduction of stomatal aperture that interacts with CO₂ diffusion as well as the intercellular CO₂ (Ci) inside the stomata; stomatal limitation effect (Cifre et al., 2005). However, at severe water deficits, Pn further declines and Ci increases indicating that non-stomatal limitations become significant (Tezara et al., 1999). Othman et al (2014a) screened leaf-level physiological changes that occurred during cyclic irrigation to determine parameters that best-represented changes in plants; the physiological variables included midday stem water potential (Ψ_{smd}), relative water content (RWC), the osmotic potential at full turgor (π), leaf area ratio (LAR), gas exchange (Pn, gs, and E) chlorophyll content and fluorescence (Fv/Fm). They found that Ψ_{smd} was the best leaf-level physiological response to detect moisture status in plants. Although the study was on woody trees, a similar trend was found in medicinal plants (Kalamartzis et al., 2020). For example, water deficit stress increased leaf temperature, Ψ_{smd} , gas exchange parameters, and dry herb yield of *Ocimum basilicum* (Kalamartzis et al., 2020).

Although physiological measurements are a reliable

source to assess plant health, the amount of time involved in data collection and the difficulties of automation of those instruments could be critical limitations to using ground physiological measurement to upscale the response to a large scale (Jones, 2004; Qarallah et al., 2021). In addition, physiological variables are at the leaf-level scale, which might be not representative of the whole tree's status (Othman et al., 2021). For example, high-performance liquid chromatography is usually used to measure pigment concentration inside the leaf, but the extraordinary cost and extraction time, along with the necessity of leaf destruction, limit its usage (Gamon and Surfus, 1999). Midday stem water potential equipment (pressure chamber) is not automated which requires the person to be onsite, requires leaf destruction, and might not be safe because of pressurized gas inside the instrument. Given that leaf-level physiological measurement is not easy, to upscale to a large scale, finding another alternative to estimate plant health is a critical issue for the plants in general and the medicinal plants' community specifically.

3. The role of remote sensing in assessing medicinal plants under drought conditions

The remote sensing approach involves the acquisition of information about an object without being in direct contact with it (Jensen, 2005). This technique enables researchers to collect information about plants in rugged topography places that are not accessible for plant-based measurements (Othman, 2014). Remote sensing is a promising approach for detecting and predicting plant morphological and physiological traits such as chlorophyll content, canopy density, and water status (Othman et al., 2014b; Santos et al., 2008). These techniques can detect, predict, and scale up leaf-level physiological responses to large areas without destroying leaves or plants (Ormeci et al., 2009). Multi-spectral satellite imagery can provide information covering large areas, while aerial photography with unmanned aerial vehicles (UAV) allows us to collect comprehensive biometric information from sites under investigation (Table 1). UAV created orthophoto with high quality that allowed confidently interprets the medicinal plants' communities during different growth stages including flowering (Fadeev et al., 2019). The combined use of multispectral aerial and satellite imagery and the high spatial resolution of UAV photography scaled up the vegetation (e.g., medicinal plants) in significant areas and accelerated the work in large areas (Fadeev et al., 2019; Qarallah et al., 2021). The age of an endangered medicinal and aromatic plant species *Valeriana jatamansi* was successfully identified using field hyperspectral remote sensing and machine learning techniques. This combined approach provides a scientific way for harvesting this plant at its optimum age avoiding its wastage (Kandpal et al., 2021). A multilevel monitoring system including spaceborne (Landsat), aerial remote sensing, and field measurement was conducted to monitor the medicinal plant (*Rheum tanguticum*) resource in Sichuan Province, China (Xie et al., 2014). They found that only the *R. tanguticum* with canopy coverages of more than 1 m² could be detected from the aerial of 10 cm resolution. Landsat alone has limited capability of detecting the scattered *R. tanguticum* plants.

Table 1. Satellite and field sensor spectral, spatial, and radiometric data and their application in vegetation studies.

Platform	Sensor	Revisit time (day)	Spatial Resolution (m)	Variable	Example
Sentinel-2A	MSI	5	10 -20	Leaf area, productivity	Cerasoli et al., (2018).
Landsat 8	OLI	16	30	Leaf area, chlorophyll, photosynthesis, water stress.	Othman et al., (2014b, 2018, 2020, 2021); Tadros et al., (2020), Sawalhah et al., (2018; 2021).
NOAA-15	AVHRR/3	1	1090	Leaf area index	Liu et al., (2012)
TERRA / AQUA	MODIS	1 – 2	250-1000	Equivalent water thickness, leaf area index, chlorophyll- <i>a</i> , fraction of photosynthetically active radiation	Cheng et al., (2013), Liu et al., (2012), Moses et al., (2009), Qu et al., (2014), Yang et al., (2006).
SPOT-5	HRS	27	5	Leaf area index and soil moisture.	Soudani et al., (2006), Gouveia et al., (2009).
	HRG	27	10		
	VEGETATION	1	165		
IKONOS-2	OSA	11- 14	4	Leaf area index, and chlorophyll- <i>a</i> .	Soudani et al., (2006), Ormeci et al., (2009).
ASD Fieldspec Pro Full Range Spectroradiometer	-		≈ 1.0 m ² at 1m nadir view (25° field-of-view)	Relative water content, chlorophyll florescent, photosynthesis, and leaf water content.	Matsushita et al., (2010), Othman et al., (2015, 2020).

The water content of plants is an essential variable for the growth and productivity of medicinal plants. The water status of plants can be monitored with surface reflectance data. Generally, canopy reflectance at near infrared (NIR, 700 – 1200 nm) is higher than at shortwave infrared (SWIR; 1300 – 2500 nm) (Figure 1). This is because leaf water status has a dominant influence in the SWIR wavelengths by strongly absorbing spectral reflectance in this spectra region (Pu et al., 2003; Eitel et al., 2006). Water status (drought) indices are commonly used to detect the potential risk of occurrence and severity of drought, and to study spatial-temporal reasoning. Remotely-sensing indices such as water band index (WBI) and band ratio (NIR/SWIR) response to water status in the plant by detecting the canopy surface reflectance in the “water bands” spectrum (1200 to 2500 nm spectral range) during drought and recovery cycles (Claudio et al., 2006; Othman et al., 2014b). Under drought conditions, the spectral reflectance in the SWIR region (water absorption region) increased and change the vegetation values of indices that rely on these bands (Eitel et al., 2006). Up to now, several remote sensing methods have been established and used for agricultural drought monitoring such as vegetation indices, band ratio and empirical remote sensing algorithms (Ligi et al., 2017; Zargar et al., 2011). These methods showed the potential results in terms of detecting drought conditions in plants (Faragó et al., 1993; Zargar et al., 2011).

The use of remote sensing techniques to monitor drought and assess its influence has been widely adopted recently (Manesh et al., 2019). It is possible to observe the intensity of plant water stress and to detect the damage in crop areas using various agricultural drought indices (Ryu et al., 2019).

This directly affects agricultural production (agricultural drought), which is most often visible in the physiological condition of plants and can be assessed from ground or satellite sensors (Wu et al., 2015; Othman et al., 2015, 2021). Using these land-surface properties, numerous satellite-based water stress indices (drought) have been developed for effective monitoring (AghaKouchak et al., 2015). However, the correlations among this vegetation are not always accurate because the definitions of indices are conceptually different based on the indicated drought phenomenon and the time scale for observing the progress (Zhang et al., 2017). In addition, the parameters used in the calculation of each index differ according to the meteorological, agricultural, and hydrological drought concepts (Anggraini et al., 2016). For example, while meteorological drought considers only atmospheric dry conditions, agricultural, and hydrological droughts are highly related to land conditions. In Iași region/Romania, the land of several aromatic and medicinal plants including *Verbena Officinalis*, *Macarof* and *Stătescu* (2017) found that Normalized Vegetation Supply Water Index (NVSWI) derived from Landsat 8 Operational Land Imager (OLI) were able to detect drought stress. Karnaris and Asimopoulos (2020) reviewed the use of the unmanned aerial vehicle for detecting aromatic plant growth change and response to harsh micro-climate in Greece. They find that UAV were able to detect vegetation cover changes (NDVI). However, more studies and vegetation indices should be developed for the Western Macedonia region to successfully detect the response of the aromatic-medicinal plants to harsh conditions (cold-dry).




Figure 1. Canopy and soil surface reflectance under well-water and drought stress conditions for woody plants. Source: Canopy reflectance data were from Bayat et al. (2016) and soil data were from Othman (2014).

Leaf pigments including chlorophyll and carotenoids are essential for several plant processes such as carbon assimilation. Chlorophyll is a green pigment found in chloroplast and present in green algae and terrestrial plants in two forms, chlorophyll, a and b. This pigment absorbs light mainly in the red (650 – 700 nm) and the blue (400 – 500 nm) and reflects the green light (~550 nm) (Jones and Vaughan, 2010). Remotely sensed data of leaf pigments especially chlorophyll can predict the leaf's physiological status and consequently the plants' health (Gamon and Surfus, 1999). Surface reflectance at 510, 550, 700 and 750 nm significantly correlated ($R^2 > 0.75$) with the total carotenoids in the leaves (Gitelson et al., 2002). Othman et al. (2018) found that multispectral data from Landsat ETM+ are a reliable source to detect chlorophyll content in the plant. In addition, Ormeci et al. (2009) concluded that IKONOS data are a reliable source for detecting chlorophyll-a in large areas even if the in-situ measurements are limited. At drought stress, excess light energy migrates from the chlorophyll molecules to the xanthophyll cycle to protect the system from damage (Grace et al., 2007). This energy causes a shift in the xanthophyll cycle, at which point violaxanthin is converted into zeaxanthin causing excess energy dissipation (Naumann et al., 2009). The hyperspectral vegetation index, and photochemical reflectance index (PRI) successfully measured the efficiency of the xanthophyll (Naumann et al., 2009). Suárez et al. (2008) found that canopy-PRI derived from Airborne Hyperspectral Scanner (AHS) sensor was able to detect the physiological responses (xanthophyll pigment cycle, stomatal conductance, and stem water potential) of plants (Suárez et al., 2008). In addition, Thenot et al. (2002) revealed that PRI could be a non-destructive, cost-effective method for detecting water stress in *Chenopodium quinoa*. Considering the previous studies that showed a significant relationship between remotely sensed data and leaf pigments, especially chlorophyll (content and fluorescence), we believe that remote sensing data are a reliable source to assess plant chlorophyll and hence plant health.

Precision agriculture of woody and herbaceous crops including medicinal plants requires a piece of accurate information about the canopy, specifically coverage percentage. Leaf area index (LAI) is an essential variable that is used to estimate vegetation cover and productivity

and as an input to water and energy budgets and ecosystem process models (Butson et al., 2002; Fernandes et al., 2004). When LAI strongly correlates with remotely sensed vegetation indices, these indices can be used to scale up those variables over large regions (Treitz et al., 2010). Othman and St. Hilaire (2021) found that three vegetation indices derived from Landsat ETM+ were able to estimate LAI correctly. The vegetation indices were normalized difference infrared Index-band 5 (NDII5), enhanced vegetation index (EVI), and green normalized difference vegetation index (GNDVI). In addition, Liu et al. (2012) concluded that vegetation indices from Landsat TM/ETM+ including normalized difference vegetation index (NDVI), the optimized soil adjusted vegetation index (OSAVI), the two bands enhanced vegetation index (EVI2) and the modified triangular vegetation index (MTVI2) can be used to derive LAI map for seasonal crop growth monitoring. Considering the finding of previous studies, the use of datasets from high-resolution aerial sensors and moderate satellites images are holding promises for detecting medicinal plants' health status by estimating canopy cover and chlorophyll content (red, 600-700 nm; near-infrared spectrum, 700-1100 nm) as well as water status (shortwave infrared, 1300-2500 nm).

4. Remote sensing limitations

The current limitations for using remotely sensed data are mainly due to restricted spectral range, coarse spatial resolution (more than 30 m), low temporal resolution (revisit time) as well as inadequate repeat coverage during the growing season (Moran et al., 1997). In addition, image pre-processing of aircraft- and satellite-based images required specialized software and workers. For example, satellite sensor data requires atmospheric and geometric correction before utilization using special software such as ENVI and ERDAS IMAGINE (Othman et al., 2018; Sawalhah et al., 2018, 2021). In addition, the acquisition of cloud-free space-born images (e.g., MODIS, Landsat) is one of the biggest challenges (Whitcraft et al., 2015). During the satellite overpass, the area should be cloud-free to guarantee meaningful images. Therefore, this tool could be inefficient during winter and early spring; the time when clouds cover percentage is extremely high. Although multispectral remotely sensed data such as Landsat series and Sentinel-2, hyperspectral remote sensing equipment is extremely expensive. In terms of physiological assessment, data from several remote sensing studies could show a pattern of difficulties in predicting or detecting the plant response. For example, when plants are under moderate water stress the difference in reflectance is quite narrow. As a result, the plant could be exposed to water stress through the surface reflectance is almost similar. At the field scale, both high spatial and high temporal data are required due to the small size of agricultural fields and the quick changes in plants through the growing season (Becker-Reshef et al., 2010; Rocha et al., 2012; Atzberger, 2013). For example, moderate spatial resolution data (i.e., 30 m) is essential for studying plant responses at a field scale (Roy et al., 2014), and high temporal resolution data (i.e., weekly) is obligatory for monitoring quick changes during the growing season (Zhang et al., 2003; Kovalsky et al., 2012). These variations,

in some cases, may reflect specific agricultural difficulties such as drought (McVicar and Jupp, 1998). Generally, if the spatial resolution is high enough (e.g., less, or equal to 30 m), then it is reasonably easy to compare with ground-based measurements. When the coarse spatial resolution is used, a combined airborne and space-borne remote sensing datasets might be used for favorable accuracy (Hazaymeh and Hassan, 2016).

5. Conclusions

Plant-based measurements including midday stem water potential, relative water content, gas exchange (Pn, gs, E and VPD), and leaf pigments (chlorophyll and carotenoids) are the best physiological measurements to assess the response of plants to environmental stresses including those planted for medicinal usage. However, those measurements are time consuming and expensive. Remotely sensed data from hyperspectral and multispectral sensors make it possible to assess medicinal plant physiology through successful detection of the ground leaf and canopy physiological variables including water status, chlorophyll, and LAI. Shortwave infrared indices such as vegetation water stress index are useful for estimating medicinal plant water status especially when ground physiological measurements (e.g., midday stem water potential) are limited. However, those remotely sensed indices can markedly predict water levels in medicinal plants under severe water stress conditions. Surface reflectance vegetation indices can be used in estimating water status in vegetation including medicinal plants when rainfall is the only source of water and when plants are exposed to severe water stress. Under commercial production of medicinal plants, the use of those indices could be not accurate because growers will not allow them to fully dry, exposing plants to severe water stress. Despite being inefficient occasionally, remote sensing sensors are a viable and accurate tool; and hold promise for assessing vegetation health including medicinal plants.

References

- Abdelhalim, A., Aburjai, T., Hanrahan, J., Abdel-Halim, H. (2017). Medicinal plants used by traditional healers in Jordan, the Tafila region. *Pharmacognosy Magazine* 13: S95-S101. doi:10.4103/0973-1296.203975.
- AghaKouchak, A., Farahmand, A., Melton, F.S., Teixeira, J., Anderson, M.C., Wardlow, B.D., Hain, C.R. (2015). Remote sensing of drought: Progress, challenges and opportunities. *Reviews of Geophysics* 53: 452–480.
- Al-Ajlouni, M., Ayad, J., Othman, Y. (2017). Increasing nutrient levels promote growth and flower quality in lilies grown under soilless culture. *Horticultural Science* 44(4):171-177.
- Al-Bakri, J., Al-Eisawi, D., Damhoureyeh, S., Oran, S. (2011). GIS-Based analysis of spatial distribution of medicinal and herbal plants in arid and semi-arid zones in the North-west of Jordan. *Annals of Arid Zone* 50(2): 99-115.
- Al-Karaki, G. and Othman, Y. (2009). Soilless cultivation of some medicinal and aromatic herb plants under the conditions of Arabian Gulf region. *Emirate Journal of Food Agriculture* 21 (2): 64-70. 3.
- Al-Kofahi, S., Gharaibeh, A., Bsoul, E., Othman, Y., St. Hilaire, R. (2019). Investigating domestic gardens' densities, spatial distribution and types among city districts. *Urban Ecosystems* 22(2). <https://doi.org/10.1007/s11252-019-0833-7>.
- Alsmirat, N., Al-Ajlouni, M., Ayad, J., Othman, Y., St. Hilaire, R. (2018). Composition of soilless substrates affect the physiology and fruit quality of two strawberry (*Fragaria × ananassa* Duch.) cultivars. *Journal of Plant Nutrition* 41(18): 2356-2364.
- Anggraini, N., Faridah, E., Indrioko, S. (2016). The effect of drought stress on physiological behavior and growth of black locust seedlings (*Robinia pseudoacacia*). *Journal of Forest Science* 9 (1): 40-56 (17).
- Arunyanark, A., Jogloy, S., Akkasaeng, C., Vorasoot, N., Kesmala, T., Rao, R., Wright, G., Patanothai, A. (2008). Chlorophyll stability is an indicator of drought tolerance in peanut. *Journal of Agronomy and Crop Science* 194: 113–125.
- A'saf, T., Al-Ajlouni, M., Ayad, J., Othman, Y., St. Hilaire, R. (2020). Performance of six different soilless green roof substrates for the Mediterranean region. *Science of the Total Environment* 730:139182. <https://doi.org/10.1016/j.scitotenv.2020.139182>.
- Atta, A.H. and Alkofahi, A. (1998). Anti-nociceptive and anti-inflammatory effects of some Jordanian medicinal plant extracts. *Journal of Ethnopharmacology* 60:117-124.
- Atzberger, C. (2013). Advances in remote sensing of agriculture: Context description, existing operational monitoring systems and major information needs. *Remote Sensing* 5: 949-981.
- Ayad, J., Othman, Y., Al-Ajlouni, M., Alsmirat, N. (2018). Photosynthesis, gas exchange and yield of two strawberry (*Fragaria × ananassa* Duch.) cultivars in response to gibberellic acid. *Fresenius Environmental Bulletin* 27: 9127-9134.
- Bayat, B.; Van der Tol, C.; Verhoef, W. (2016). Remote sensing of grass response to drought stress using spectroscopic techniques and canopy reflectance model inversion. *Remote Sensing* 8, 557. <https://doi.org/10.3390/rs8070557>.
- Becker-Reshef, I., Justice, C., Sullivan, M., et al. (2010). Monitoring global croplands with coarse resolution earth observations: The global agriculture monitoring (GLAM) project. *Remote Sensing* 2: 1589-1609.
- Bhuiyan, C., Singh, R. P., Kogan, F. N. (2006). Monitoring drought dynamics in the Aravalli region (India) using different indices based on ground and remote sensing data. *International Journal of Applied Earth Observation and Geoinformation* 8:289–302.
- Blackburn, G. (2007). Hyperspectral remote sensing of plant pigments. *Journal of Experimental Botany* 58(4): 855-867.
- Blum, A. (2017). Osmotic adjustment is a prime drought stress adaptive engine in support of plant production. *Plant, Cell and Environment* 40: 4–10.
- Butson, C., Fernandes, R., Latifovic, R., Wenjun, C. (2002). A robust approach for estimating LAI from Landsat TM/ETM + imagery. *Proceedings of the IEEE Int. Geosciences Remote Sensing* 4: 2328–2330. DOI: 10.1109/IGARSS.2002.1026534.
- Cerasoli, S., Campagnolo, M., Faria, J., Nogueira, C., Caldeira, M. (2018). On estimating the gross primary productivity of Mediterranean grasslands under different fertilization regimes using vegetation indices and hyperspectral reflectance. *Biogeosciences* 15: 5455–5471.
- Chapman, K. and Chomchalow, N. (2005). Production of medicinal plants in Asia. *Acta Horticulturae*: 679: 45–59.
- Cheng, T., Riano, D., Koltunov, A., Whiting, M., Ustin, S., Rodriguez, J. (2013). Detection of diurnal variation in orchard canopy water content using MODIS/ASTER airborne simulator (MASTER) data. *Remote Sensing of Environment* 132:1–12.
- Cifre, J., Bota, J., Escalona, J., Medrano, H., Flexas, J. (2005). Physiological tools for irrigation scheduling in grapevine (*Vitis vinifera* L.) An open gate to improve water-use efficiency. *Agriculture, Ecosystems and Environment* 106:159–170.
- Claudio, H., Cheng, Y., Fuentes, D., Gamon, J., Luo, H., Oechel, W., Qiu, H., Rahman, A., Sims, D. (2006). Monitoring drought effects on vegetation water content and fluxes in chaparral with

- the 970 nm water band index. *Remote Sensing of Environment* 103:304–311.
- Damour, G., Vandame, M., Urban, L. (2009). Long-term drought results in a reversible decline in photosynthetic capacity in mango leaves, not just a decrease in stomatal conductance. *Tree Physiology* 29: 675–684.
- Eitel, J., Gessler, P., Smith, A., Robberecht, R. (2006). Suitability of existing and novel spectral indices to remotely detect water stress in *Populus* spp. *Forest Ecology and Management* 229:170–182.
- Fadeev, N., Skrypitsyna, T., Kurkov, V., Sidelnikov, N. (2019). Use of remote sensing data and GIS technologies for monitoring stocks of medicinal plants: Problems and Prospects. *Information Technologies in the Research of Biodiversity, SPEES* 4-21.
- FAO (Food and Agriculture Organization, 2008). Climate change, water, and food security. Land and Water Division.
- Faragó, T., Kozma, E., Nemes, Cs. (1993). Drought indices in meteorology. *Quarterly Journal of the Hungarian Meteorological Service* 93(1): 45–60.
- Fernandes, R. A., Miller, J. R., Chen, J. M., Rubinstein, I. G. (2004). Evaluating image-based estimates of leaf area index in boreal conifer stands over a range of scales using high-resolution CASI imagery. *Remote Sensing of Environment* 89: 200-216. <https://doi.org/10.1016/j.rse.2002.06.005>.
- Gamon, J. and Surfus, J. (1999). Assessing leaf pigment content and activity with a reflectometer. *New Phytology* 143(1): 105-117.
- García-Caparrós, P., Romero, M.J., Llanderal, A., Cermeño, P., Lao, M.T., Segura, M.L. (2019). Effects of drought stress on biomass, essential oil content, nutritional parameters, and costs of production in six Lamiaceae species. *Water* 11: 573. <https://doi.org/10.3390/w11030573>.
- Gitelson, A., Zur, Y., Chivkunova, O., Merzlyak, M. (2002). Assessing carotenoid content in plant leaves with reflectance spectroscopy. *Photochemistry and Photobiology* 75(3): 272-281.
- Gori, A., Brunetti, C., dos Santos Nascimento, L.B., Marino, G., Guidi, L., Ferrini, F., Centritto, M., Fini, A., Tattini, M. (2021). Photoprotective role of photosynthetic and non-photosynthetic pigments in *Phillyrea latifolia*: Is their “Antioxidant” function prominent in leaves exposed to severe summer drought? *International Journal of Molecular Sciences* 22: 8303, <https://doi.org/10.3390/ijms22158303>.
- Gouveia, C., Trigo, R., Dacamara, C. (2009). Drought and vegetation stress monitoring in Portugal using satellite data. *Natural Hazards and Earth System Sciences* 9:185–195.
- Grace, J., Nichol, C., Disney, M., Lewis, P., Quaife, T., Bowyer, P. (2007). Can we measure terrestrial photosynthesis from space directly, using spectral reflectance and fluorescence? *Global Change Biology* 13(7): 1484-1497.
- Grams, T., Koziol, C., Lautner, S., Matyssek, R., Fromm, J. (2007). Distinct roles of electric and hydraulic signals on the reaction of leaf gas exchange upon re-irrigation in *Zea mays* L. *Plant, Cell and Environment* 30(1): 79-84.
- Guerfel, M., Baccouri, O., Boujnah, D., Zarrouk, M. (2008). Changes in lipid composition, water relations and gas exchange in leaves of two young ‘Chmlali’ and ‘Chetoui’ olive trees in response to water stress. *Plant Soil* 311: 121–129.
- Guidi, L., Lo Piccolo, E., Landi, M. (2019). Chlorophyll fluorescence, photoinhibition and abiotic stress: does it make any difference the fact to be a C3 or C4 species? *Front. Plant Science* 10:174. doi: 10.3389/fpls.2019.00174.
- Hao, D. (2019). Genomics and Evolution of Medicinal Plants. In: *Ranunculales Medicinal Plants Biodiversity, Chemodiversity and Pharmacotherapy* pp 1-33.
- Hazaymeh, K. and Hassan, Q. K. (2017). A remote sensing-based agricultural drought indicator and its implementation over a semi-arid region, Jordan. *Journal of Arid Land* 9:319–330.
- Hazaymeh, K. and Hassan, Q.K. (2016). Remote sensing of agricultural drought monitoring: A state of art review. *AIMS Environmental Science* 3(4): 604-630. doi: 10.3934/environsci.2016.4.604.
- He, C., Zhang, J., Duan, A., Sun, H., Fu, L., Zheng, S. (2007). Proteins responding to drought and high temperature stress in *Pinus armandii* Franch. *Canadian Journal for Research* 85: 994-1002.
- IPCC. (2013). Intergovernmental Panel on Climate Change. Climate change: the physical science basis. In: Stocker, T.F., Qin, D., Plattner, M., Tignor, S.K., Allen, et al. (eds.), Working group I contribution to the fifth assessment report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Jeelani, S.M., Rather, G.A., Sharma, A., Lattoo, S.K. (2018). In perspective: Potential medicinal plant resources of Kashmir Himalayas, their domestication and cultivation for commercial exploitation. *Journal of Applied Research on Medicinal and Aromatic Plants* 8:10–25.
- Jensen, J. (2005). Introductory digital image processing: A remote sensing perspective. Pearson Prentice Hall. Third edition. USA.
- Jones, H. (2004). Irrigation scheduling: Advantages and pitfalls of plant-based methods. *Journal of Experimental Botany* 55:2427–2436.
- Jones, H. and Vaughan, R. (2010). Remote sensing of vegetation, principles, techniques, and applications. Oxford University Press. USA.
- Kalamartzis, I., Menexes, G., Georgiou, P., Dordas, C. (2020). Effect of water stress on the physiological characteristics of five Basil (*Ocimum basilicum* L.) Cultivars. *Agronomy* 10: 1029. <https://doi.org/10.3390/agronomy10071029>.
- Kandpal K., Kumar S., Venkat G., Meena R., Pal P., Kumar, A. (2021). Onsite age discrimination of an endangered medicinal and aromatic plant species *Valeriana jatamansi* using field hyperspectral remote sensing and machine learning techniques. *International Journal of Remote Sensing* 42(10): 3777-3796, DOI: 10.1080/01431161.2021.1881184.
- Karnaris, I., and Asimopoulos, N. (2020). Overview of Aromatic Plants Precision Agriculture with the use of UAV. Proceeding of the 9th International Conference on Information and Communication Technologies, Agriculture, Food and Environment (HAICTA, 2020), Thessaloniki, Greece, pages 24-27.
- Khasawneh, A., Alsmairat, N., Othman, Y., Ayad, J., Al-Qudah T., Leskovar, D. (2021). Influence of nitrogen source on physiology, yield and fruit quality of young apricot trees. *Journal of Plant Nutrition* <https://doi.org/10.1080/01904167.2021.1918718>.
- Khorasaninejad, S., Mousavi, A., Soltanloo, H., Hemmati, K., Khalighi, A. (2010). The effect of salinity stress on growth parameters, essential oil yield and constituent of peppermint (*Mentha piperita* L.). *World Applied Sciences Journal* 11 (11): 1403-1407.
- Kovalskyy, V., Roy, D.P., Zhang, X. Y., et al., (2012). The suitability of multi-temporal web-enabled Landsat data NDVI for phenological monitoring—a comparison with flux tower and MODIS NDVI. *Remote Sensing Letters* 3: 325-334.
- Leskovar, D. and Othman Y. (2018). Organic and conventional farming differentially influenced soil respiration, physiology, growth and head quality of artichoke cultivars. *Journal of Soil Science and Plant Nutrition* 18 (3), 865-880.
- Leskovar, D. and Othman, Y. (2016). Low nitrogen and fertigation practices can improve artichoke transplant quality and yield. *HortScience* 51(5):567-572. <https://doi.org/10.21273/HORTSCI.51.5.567>.
- Leskovar, D. and Othman, Y. (2019). Nitrogen management for

- improving root and shoot components of young 'Arbequina' Olives. *HortScience* 54(1):175-180. <https://doi.org/10.21273/HORTSCI13397-18>.
- Leskovar, D. and Othman, Y. (2021). Direct Seeding and transplanting influence root dynamics, morpho-physiology, yield, and head quality of globe artichoke. *Plants* 10, 899. <https://doi.org/10.3390/plants10050899>.
- Li, Y., He, N., Hou, J., Xu, L., Liu, C., Zhang, J., Wang, Q., Zhang, X., Wu, X. (2018). Factors influencing leaf chlorophyll content in natural forests at the biome scale. *Frontiers in Ecology and Evolution* 6:64. doi:10.3389/fevo.2018.00064.
- Ligi, M., Kutser T., Kallio K, Attila J., Koponen S., Paavel B., Soomets T., Reinart A. (2017). Testing the performance of empirical remote sensing algorithms in the Baltic Sea waters with modelled and in situ reflectance data. *Oceanologia* 59: 57-68.
- Liu, J., Pattey, E., Jégo, G. (2012). Assessment of vegetation indices for regional crop green LAI estimation from Landsat images over multiple growing seasons. *Remote Sensing of Environment* 123: 347–358. <https://doi.org/10.1016/j.rse.2012.04.002>.
- Macarof, P., and Stătescu F. (2017). Identification drought extent based on NVSWI using Landsat data: a case study of Iași County. *Lucrări Științifice* 60:221-224.
- Manesh, M.B., Khosravi, H., Alamdarloo, E. H., Alekasir, M. S., Gholami, A., Singh, V.P. (2019). Linkage of agricultural drought with meteorological drought in different climates of Iran. *Theoretical and Applied Climatology* 1–9.
- Matsushita, B., Xu, M., Onda, Y., Otsuki, Y., Toyota, M. (2010). Detecting forest degradation in Kochi, Japan: ground-based measurements versus satellite (Terra/ASTER) remote sensing. *Hydrological Processes* 24:588–595.
- McVicar, T. R. and Jupp, D. L. (1998). The current and potential operational uses of remote sensing to aid decisions on drought exceptional circumstances in Australia: A review. *Agricultural Systems* 57: 399-468.
- Moran, M., Inoue Y. and Barnes E. (1997). Opportunities and limitations for image-based remote sensing in precision crop management. *Remote Sensing of Environment* 61:319-346.
- Moses, W., Gitelson, A., Berdnikov, S., Povazhnyy, V. (2009). Estimation of chlorophyll-a concentration in case II waters using MODIS and MERIS data successes and challenges. *Environmental Research Letters* 4:1–8.
- Naumann, J., Young, D., and Anderson, J. (2009). Spatial variations in salinity stress across a coastal landscape using vegetation indices derived from hyperspectral imagery. *Plant Ecology* 202: 285-297.
- NDMC. (2021). Type of drought. National Drought Mitigation Center. University of Nebraska. Accessed, October 25, 2021, <https://drought.unl.edu/Education/DroughtIn-depth/TypesofDrought.aspx>.
- Oran S. A. and Al-Eisawi D. M. (2015). Ethnobotanical survey for the medicinal plants in the high mountains of Jordan. *Journal of Biodiversity and Environmental Sciences* 6 (3), 381-400.
- Ormeci, C., Sertel, E., Sarikaya, O. (2009). Determination of chlorophyll-a amount in golden Horn, Istanbul, Turkey using IKONOS and in situ data. *Environmental Monitoring Assessment* 155:83–90.
- Othman, Y. (2014). Detecting moisture status of pecan orchards and the potential of remotely-sensed surface reflectance data. New Mexico State University, NM, USA.
- Othman, Y. and Leskovar D. (2018). Organic soil amendments influence soil health, yield, and phytochemicals of globe artichoke heads. *Biological Agriculture and Horticulture* 34:4, 258-267. DOI:10.1080/01448765.2018.1463292.
- Othman, Y. and Leskovar, D. (2019). Nitrogen management influenced root length intensity of young olive trees. *Scientia Horticulturae* 246:726–733.
- Othman, Y. and St. Hilaire, R. (2021). Using multispectral data from Landsat ETM+ to estimate leaf area index of pecan orchards. *Fresenius Environmental Bulletin* 30:2613-2618.
- Othman, Y., Al-Ajlouni, M., A'saf, T., Sawalhah, H., Bany Hani, M. (2021). Influence of gibberellic acid on the physiology and flower quality of gerbera and lily cut flowers. *International Journal of Agriculture and Natural Resources* 48(1): 21-33, <http://dx.doi.org/10.7764/ijanr.v48i1.2218>.
- Othman, Y., Al-Ajlouni, St. Hilaire, R. (2019). Using hyperspectral surface reflectance data to detect chlorophyll content in pecans. *Fresenius Environmental Bulletin* 28(8): 6117-6124.
- Othman, Y., Steele, C., St. Hilaire, R. (2018). Surface Reflectance Climate Data Records (CDRs) is a reliable Landsat ETM+ source to study chlorophyll content in pecan orchards. *Journal of the Indian Society of Remote Sensing* 46(2): 211–218.
- Othman, Y., Steele, C., Van Leeuwen, D., St. Hilaire, R. (2015). Hyperspectral surface reflectance data used to detect moisture status of pecan orchards during flood irrigation. *Journal of the American Society for Horticultural Science* 140(5): 449-458.
- Othman, Y., Van Leeuwen, D., Heerema, R., St. Hilaire, R. (2014a). Midday stem water potential values needed to maintain photosynthesis and gas exchange established for pecans. *Journal of the American Society for Horticultural Science* 139(5): 537–546.
- Othman, Y., Steele, C., Van Leeuwen, D., Heerema, R., Bawazir, S., St. Hilaire, R. (2014b). Remote sensing used to detect moisture status of pecan orchards grown in a desert environment. *International Journal of Remote Sensing* 35(3): 949-966.
- Pavarini, D. P., Pavarini, S. P., Niehues, M., Lopes, N. P. (2012). Exogenous influences on plant secondary metabolite levels. *Animal Feed Science and Technology* 176(1-4): 5–16.
- Pirzad, A., Shakiba, M., Zehtab-Salmasi, S., Mohammadi, S., Darvishzadeh, R., Samadi, A. (2011). Effect of water stress on leaf relative water content, chlorophyll, proline and soluble carbohydrates in *Matricaria chamomilla* L. *Journal of Medicinal Plants Research* 5(12): 2483-2488.
- Pu, R., Ge, S., Kelly, N., Gong, P. (2003). Spectral absorption features as indicators of water status in coast live oak (*Quercus agrifolia*) leaves. *International Journal of Remote Sensing* 24:1799–1810.
- Qarallah, B., Al-Ajlouni, M., Al-Awasi, A., Alkarmy, M., Al-Qudah, E., Bani Naser, A., Al-Assaf, A., Gevaert, C., Al Asmar, Y., Belgiu, M., Othman, Y. (2021). Evaluating post-fire recovery of Latroon dry forest using Landsat ETM+, unmanned aerial vehicle and field survey data. *Journal of Arid Environments* 193:104587.
- Qu, Y., Zhu, Y., Han, W., Wang, J., Ma, M. (2014). Crop leaf area index observations with a wireless sensor network and its potential for validating remote sensing products. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* 7:431–443.
- Razmjoo, K. P., Heydarizadeh, Sabzalian, M. R. (2008). Effect of salinity and drought stresses on growth parameters and essential oil content of *Matricaria chamomilla*, *International Journal of Agriculture and Biology*.
- Rocha, J., Perdigão, A., Melo, R., et al. (2012). Remote sensing based crop coefficients for water management in agriculture. In: Curkovic, S. Sustainable Development—Authoritative and Leading Edge Content for Environmental Management 167-192.
- Rousta, I., Olafsson, H., Moniruzzaman, M., Zhang, H., Liou, Y., Mushore, T.D., Gupta, A. (2020) Impacts of drought on vegetation assessed by vegetation indices and meteorological factors in Afghanistan. *Remote Sensing* 12: 2433. <https://doi.org/10.3390/rs12152433>.

- Roy, D. P., Wulder, M. A., Loveland, T. R., et al. (2014). Landsat-8: Science and product vision for terrestrial global change research. *Remote Sensing of Environment* 145: 154-172.
- Ryu, J., Han, K., Lee, Y., Park, N., Hong, S., Chung, C., Cho, J. (2019). different agricultural responses to extreme drought events in neighboring counties of south and north Korea. *Remote Sensing* 11:1773. <https://doi.org/10.3390/rs11151773>.
- Santos, C., Lorite, J., Tasumi, M., Allen, R., Fereres, E. (2008). Integrating satellite-based evapotranspiration with simulation models for irrigation management at the scheme level. *Irrigation Science* 26(3):277-288.
- Sawalhah, M., Al-Kofahi, S., Othman, Y., Cibils, A. (2018). Assessing rangeland cover conversion in Jordan after the Arab spring using a remote sensing approach. *Journal of Arid Environments* 157:97-102.
- Sawalhah, M., Othman, Y., Abu Yahya, A., Al-Kofahi, S., Al-Lataifeh, F., Cibils, A. (2021). Evaluating the influence of COVID-19 pandemic lockdown on Jordan Badia rangelands. *Arid Land Research and Management*, 35: 483-495, DOI: 10.1080/15324982.2021.1921071.
- Seneviratne, S. I., Nicholls, N., Easterling, D. C., Goodess, M., Kanae, S., Kossin, J., Luo, Y., Marengo, J., McInnes, K., Rahimi, M., Reichstein, M., Sorteberg, A., Vera, C., Zhang, X. (2012). Changes in climate extremes and their impacts on the natural physical environment. Cambridge, UK, and New York, NY, USA, pp. 109-230.
- Sharma, T., Kaur, A., Saajan, S., Thakur, R. (2020). Effect of nitrogen on growth and yield of medicinal plants: A review paper. *European Journal of Molecular and Clinical Medicine* 7: 2771-2776.
- Soudani, K., Francois, C., Maire, G., Dantec, V., Dufrene, E. (2006). Comparative analysis of IKONOS, SPOT, and ETM+ data for leaf area index estimation in temperate coniferous and deciduous forest stands. *Remote Sensing of Environment* 102:161-175.
- Suárez, L., Zarco-Tejada, P. J., Sepulcre-Cantó, G., Pérez-Priego, O., Miller, J.R., Jiménez-Muñoz, J.C., Sobrino, J. (2008). Assessing canopy PRI for water stress detection with diurnal airborne imagery. *Remote Sensing of Environment* 112: 560-575.
- Tadros, M. J., Al-Mefleh, N., Othman, Y., Al-Assaf, A. (2021). Water harvesting techniques for improving soil water content, and morpho-physiology of pistachio trees under rainfed conditions. *Agricultural Water Management* 243: 106464. <https://doi.org/10.1016/j.agwat.2020.106464>.
- Tadros, M.J., Al-Assaf, A., Othman, Y.A., Makhmreh, Z., Taifour, H. (2020). Evaluating the effect of *Prosopis juliflora*, an alien invasive species, on land cover change using remote sensing approach. *Sustainability* 12:5887, <https://doi.org/10.3390/su12155887>.
- Tahat, M., Alananbeh, K., Othman, Y., Leskovar, D. (2020). Soil health and sustainable agriculture. *Sustainability* 12: 4859. <https://doi.org/10.3390/su12124859>.
- Tanga, M., Lewu, F., Oyedeji, O., Oyedeji, O. (2018). Cultivation of medicinal plants in South Africa: A solution to quality assurance and consistent availability of medicinal plant materials for commercialization. *Academia Journal of Medicinal Plants* 6: 168-177.
- Tardieu, F., Simonneau, T., Muller, B. (2018). The physiological basis of drought tolerance in crop plants: A scenario-dependent probabilistic approach. *Annual Review of Plant Biology* 69: 733-759.
- Tátrai, Z., Sanoubar, R., Pluhár, Z., Mancarella, S., Orsini, F., Gianquinto, G. (2016). Morphological and physiological plant responses to drought stress in *Thymus citriodorus*. *International Journal of Agronomy*, 4165750: 8 <https://doi.org/10.1155/2016/4165750>.
- Tezara, W., Mitchell, V., Driscoll, S., Lawlor, D. (1999). Water stress inhibits plant photosynthesis by decreasing coupling factor and ATP. *Nature* 401: 914-917.
- Thenot, F., Methy, M., Winkel, T. (2002). The Photochemical Reflectance Index (PRI) as a water-stress index. *International Journal of Remote Sensing* 23(23): 5135-5139.
- Treitz, P., Thomas, V., Zarco-Tejada, P., Gong, P., Curran, P. (2010). Hyperspectral remote sensing for forestry. *American Society for Photogrammetry and Remote Sensing (ASPRS)*. USA.
- Trenberth, K.E., Dai, A., van der Schrier, G., Jones, P. D., Barichivich, J., Briffa, K. R., Sheffield, J. (2014). Global warming and changes in drought. *Nature Climate Change* 4: 17-22. DOI: 10.1038/nclimate2067.
- UNDRR, 2021. United Nations Office for Disaster Risk Reduction. GAR Special Report on Drought 2021. Geneva. Accessed October 26, 2021, <https://www.undrr.org/publication/gar-special-report-drought-2021>.
- Van Lanen and Peters, E. (2000). Definition, Effects and Assessment of Groundwater Droughts. In Vogt, J., Somma, F. Drought and drought mitigation in Europe. Pages 49-61.
- Warren, C., Bleby, T., Adams, M. (2007). Changes in gas exchange versus leaf solutes as a means to cope with summer drought in *Eucalyptus marginata*. *Oecologia* 154:1-10.
- WB. 2021. The world bank. climate-smart agriculture. accessed September 3, 2021. <https://www.worldbank.org/en/topic/climate-smart-agriculture>.
- Whitcraft A.K., Vermote E.F., Becker-Reshef I., Justice C.O. (2015). Cloud cover throughout the agricultural growing season: impacts on passive optical earth observations. *Remote Sensing of Environment* 156: 438-447.
- Wilhite, D.A., Glantz, M. (1985). Understanding the drought phenomenon: The role of definitions. *Water International* 10(3):111-120.
- Wu, D., Qu, J. J., Hao, X. (2015). Agricultural drought monitoring using MODIS-based drought indices over the USA Corn Belt. *International Journal of Remote Sensing* 36: 5403-5425.
- Xie, C., Song, J., Suo, F., Li, X., Li, Y., Yu, H., Xu, X., Luo, K., Li, Q., Xin, T., Guan, M., Xu, X., Miki, E., Takeda, Q., Chen, S. (2014). Natural resource monitoring of *Rheum tanguticum* by multilevel remote sensing. *Evidence-Based Complementary and Alternative Medicine* 618902: 9 <https://doi.org/10.1155/2014/618902>.
- Yang, W., Tan, B., Huang, D., Rautiainen, M., Shabanov, N., Wang, Y., Privette, J., Huemmrich, K., Fensholt, R., Sandholt, I., Weiss, M., Ahl, D., Gower, S., Nemani, R., Knyazikhin, Y., Myneni, R. (2006). MODIS leaf area index products: from validation to algorithm improvement. *IEEE Transactions on Geosciences and Remote Sensing* 44:1885-1897.
- Yurekli, K. and Kurunc, A. (2004). Simulation of drought periods using stochastic models. *Turkish Journal of Engineering and Environmental Sciences* 28:181-190.
- Zaid, A., Ahmad, B., Wani, S.H. (2021) Medicinal and Aromatic Plants Under Abiotic Stress: A Crosstalk on Phytohormones' Perspective. In: Aftab T., Hakeem K.R. (eds) *Plant Growth Regulators*. Springer, Cham. https://doi.org/10.1007/978-3-030-61153-8_5.
- Zargar, A., Sadiq, R., Naser, B., Khan, F. I. (2011). A review of drought indices. *Environmental Reviews* 19: 333-349. DOI: 10.1139/a11-013.
- Zhang, X., Friedl, M. A., Schaaf, C. B., et al. (2003). Monitoring vegetation phenology using MODIS. *Remote Sensing of Environment* 84: 471-475.
- Zhang, X., Wu, S., Yan, X., Chen, Z. (2017). A global classification of vegetation based on NDVI, rainfall and temperature. *International Journal of Climatology* 37: 2318-2324.