

---

**PAGES****PAPERS**

---

- 99 - 105 Effectiveness of water harvesting technique on selected pastoral shrub: a case of the Syrian Badia (Qaryatien)  
*Mahmoud Al Hamdan, Wafaa Aboud, Boshra Khozam, Issam AlKhouri*
- 106 - 112 Source rock evaluation of the Chia Gara Formation in the Bekhme-1 well, Harir District, Kurdistan Region, Iraq  
*Rzger Abdula, Sardar Fatah, Gardun Salih, Mohammed Mustafa, Muhammed Ali*
- 113- 121 Depositional environment, microfacies analysis and planktonic foraminifera of oil shale deposits in the Wadi Al-Shallala Area, NW Jordan.  
*Hanan Al-Soud and Habes Al-Mashakbeh*
- 122 - 133 Treatment of real olive mill wastewater by sole and combination of  $H_2O_2$ ,  $O_3$ , and UVA: effect of doses and ratios on organic content and biodegradability  
*Dheaya Alrousan*
- 134 - 144 Analysis of socio-economic and housing characteristics in some selected slum area in Lagos State Metropolis, Nigeria using Geographical Information System  
*Oluwaseun Okimiji, Oludare Adedeji, Olusegun Oguntoke, Olufunke Shittu, Moses Aborisade, Obioma Ezennia1*
- 145 - 153 Architectural design solutions for combating dust storms in residential buildings (case study: Abadan City, Iran)  
*Amena Agharabi and Zeinab Fard*
- 154 - 162 A designed model for identifications of *Dicarinella concavata* (Brotzen, 1934) and *Dicarinella asymmetrica* (Sigal, 1952) planktic foraminifer species under thin sections: an example from the Kurdistan region, NE Iraq  
*Rawand Bakir Noori Jaff*
- 163 - 186 Paleocene rotaliid benthic foraminifera of Jabal Mundassa, Al Ain area, United Arab Emirates  
*Haidar Salim Anan*
- 
-